

**Gene
Synthesis**

**DNA
Template**

**RNA
Synthesis**


**Primer
pools**

Other


Oligonucleotide Library



Cleavage chemistry developed to release ligo-nucleotides from solid support with free 3'-OH



PGR Chemistry



Parallel microfluidic synthesizer chip



Digital Photolithography

Figure 1




Figure 2




Figure 3

10/533208




Figure 4

10/533208




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9

Fusion PCR for Multi-Stage Long Gene Assembling




Figure 10




Figure 11




Figure 12




Figure 13





Figure 14



1. In situ synthesized oligos (about 60mer). Each oligo contains a rU, one T7 promoter, specific RNAi sequence and a Mly I restriction site. Mly I is a blunt cutting enzyme (GAGTCNNNNN⁺).
2. Oligos are cleaved by RNase A or chemicals.
3. Flash the oligos into tube. PCR amplify the targets with universal primers, one contains the T7 sequence and the other contains the restriction sequence.
4. Double stranded PCR products with thousands of specific RNAi sequences and common T7 and restriction site.
5. Digest the PCR products with Mly I enzyme to remove the tail.
6. In vitro transcription to produce unlimited supply of RNAi pools (can be re-amplified and re-transcribed) ready for use.

Figure 15




Figure 16




Figure 17





Figure 18



Dish 1

Dish 2

Figure 19



Figure 20




Figure 21




Figure 22




Figure 23





Figure 24

Synthesis of DNA Oligos Greater than 100 bases



100-mer synthesis
validated by hybridization
(99% stepwise yield)

Figure 25




Figure 26




Figure 27




Figure 28






Figure 29

Electrophoresis
gel analysis of
PCR sequences
(60 mers) and the
sequences after
SAP1 enzyme
digestion (41 and
19 mers)



Perfect
Match
Single Deletion
Double Deletion

A



B

Probes
Figure 30

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.